


panTRK, RMab**Clone: RBT-TRK**

Rabbit Monoclonal

**RUO**
Bio SB
 BIOSCIENCE FOR THE WORLD

www.biosb.com

Inset: IHC panTRK on a Papillary Thyroid Carcinoma Tissue

Intended Use

For Research Use Only.

This antibody is intended for use in Immunohistochemical applications on formalin-fixed paraffin-embedded tissues (FFPE), frozen tissue sections and cell preparations. Interpretation of results should be performed by a qualified medical professional.

Immunogen

Synthetic peptide corresponding to the C-terminal residues of human pan TRK protein.

Summary and Explanation

TRK (tropomyosin receptor kinase) A, B, and C are encoded by *NTRK1*, *NTRK2*, and *NTRK3* genes respectively. Each protein is activated by different neurotrophins: TRKA is activated by Nerve growth factor, TRKB by brain-derived neurotrophic factor, and TRKC by NT-3. The TRK receptors are a family of tyrosine kinases that regulate synaptic strength and plasticity in the mammalian nervous system. The activation of TRK receptors by neurotrophin binding may lead to activation of signal cascades resulting in promoting survival and other functional regulation of cells. TRK family of receptor tyrosine kinases are of interest as the *NTRK* genes that encode them are involved in gene fusions identified in a wide range of adult and pediatric tumors. TRK, was initially identified in a colon carcinoma, is frequently activated in thyroid papillary carcinomas.

Pan-TRK IHC has shown to be positive in most cases with *NTRK* fusion transcripts confirmed by Archer. One study established the Pan-TRK IHC sensitivity and specificity for transcribed *NTRK* fusions to be 95.2% and 100%, respectively. All positive IHC cases had cytoplasmic staining while the following fusion partner-specific patterns were discovered: all *LMNA-NTRK1* fusions displayed nuclear membrane accentuation, all *TPM3/4* fusions displayed cellular membrane accentuation, and half of *ETV6-NTRK3* fusions displayed nuclear staining. In another study, Immunohistochemistry screening in 1043 various solid tumors showed TRKA expression in 1.6% of samples, including Colorectal, Lung Cancer, Biliary Carcinoma and Thyroid Cancers. *NTRK* gene fusions have been identified in both pediatric and adult primary central nervous system tumors, including Glioblastoma Multiforme, Pediatric Gliomas and Astrocytomas. Various translocations involving *NTRK1* or *NTRK3* have been reported in Spitzoid melanocytic neoplasms, as well as in compound Spitz Nevi. TRK fusions have also been reported in Intrahepatic Cholangiocarcinomas, Breast Cancer, quadruple wild-type (ETV6-NTRK3) Gastrointestinal Stromal Tumors, Gallbladder Adenocarcinomas, Pancreatic Carcinomas, Sinus-Nasal Low-Grade Non-Intestinal-type Adenocarcinomas and Neuroendocrine Tumors of the small bowel. In addition to being present in solid tumors, *NTRK* gene fusions have been also detected in Acute Lymphoblastic Leukemia and Acute Myeloid Leukemia.

Antibody Type	Rabbit Monoclonal	Clone	RBT-TRK
Isotype	IgG	Reactivity	Paraffin, Frozen
Localization	Cytoplasmic, Nuclear	Control	Brain, Lung Neuroendocrine
Species Reactivity		Human	

Presentation

Anti-pan TRK is a rabbit monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH 7.5, containing BSA and sodium azide as a preservative.

Catalog No.	Antibody Type	Dilution	Volume/Qty
BSB-2376-3	Tinto Predilute	Ready-to-Use	3.0 mL
BSB-2376-7	Tinto Predilute	Ready-to-Use	7.0 mL
BSB-2376-15	Tinto Predilute	Ready-to-Use	15.0 mL
BSB-2376-01	Concentrate	1:10 - 1:50	0.1 mL
BSB-2376-05	Concentrate	1:10 - 1:50	0.5 mL
BSB-2376-1	Concentrate	1:10 - 1:50	1.0 mL

Control Slides Available

Catalog No.	Quantity
BSB-2376-CS	5 slides

Precautions

1. For professional users only. Results should be interpreted by a qualified medical professional.
2. This product contains <0.1% sodium azide (NaN₃) as a preservative. Ensure proper handling procedures are used with this reagent.
3. Always wear personal protective equipment such as laboratory coat, goggles and gloves when handling reagents.
4. Dispose of unused solution with copious amount of water.
5. Do not ingest reagent. If reagent is ingested, seek medical advice immediately.
6. Avoid contact with eyes. If contact occurs, flush with large quantities of water.
7. Follow safety precautions of the heating device used for epitope retrieval (TintoRetriever Pressure Cooker or similar).
8. For additional safety information refer to Safety Data Sheet for this product.
9. For complete recommendations for handling biological specimens, please refer to the CDC document, "Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories" (see References in this document).

Storage Store at 2-8°C

Stability

This product is stable up to the expiration date on the product label. Do not use after expiration date listed on package label. Temperature fluctuations should be avoided. Store appropriately when not in use and avoid prolonged exposure to room temperature conditions.

Specimen Preparation

Paraffin sections: The antibody can be used on formalin-fixed paraffin-embedded (FFPE) tissue sections. Ensure tissue undergoes appropriate fixation for best results. Pre-treatment of tissues with heat-induced epitope retrieval (HIER) is recommended using Bio SB ImmunoDNA Retriever with Citrate (BSB 0020-BSB 0023), ImmunoDNA Retriever with EDTA (BSB 0030-BSB 0033) or ImmunoDNA Digestor (BSB 0108-0112). See reverse side for complete protocol. Tissue should remain hydrated via use of Bio SB Immuno/DNA Washer solutions (BSB 0029 & BSB 0042).

Frozen sections and cell preparations: The antibody can be used for labeling acetone-fixed frozen sections and acetone-fixed cell preparations.

Staining Procedure

1. Cut and mount 3-5 micron formalin-fixed paraffin-embedded tissues on positively charged slides such as Bio SB Hydrophilic Plus Slides (BSB 7028).
2. Air dry for 2 hours at 58° C.
3. Deparaffinize, dehydrate and rehydrate tissues.
4. Subject tissues to heat induced epitope retrieval (HIER) using a suitable retrieval solution such as ImmunoDNA Retriever with Citrate (BSB 0020-BSB 0023) or EDTA (BSB 0030-BSB 0033).
5. Any of three heating methods may be used:

a. TintoRetriever Pressure Cooker or Equivalent

Place tissues/slides in a staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA, and place on trivet in the pressure cooker. Add 1-2 inches of distilled water to the pressure cooker and turn heat to high. Incubate for 15 minutes. Open and immediately transfer slides to room temperature.

b. TintoRetriever PT Module or Water Bath Method

Place tissues/slides in a pre-warmed staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA at 95°-99° C. Incubate for 30-60 minutes.

c. Conventional Steamer Method

Place tissues/slides in a pre-warmed staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA in a steamer, cover and steam for 30-60 minutes.

6. After heat treatment, transfer slides in ImmunoDNA Retriever with Citrate or EDTA to room temperature and let stand for 15-20 minutes.
7. For manual staining, perform antibody incubation at ambient temperature. For automated staining methods, perform antibody incubation according to instrument manufacturer's instructions.
8. Wash slides with ImmunoDNA washer or DI water.
9. Continue IHC staining protocol. Wash slides between each step with ImmunoDNA washer solution.

Abbreviated Immunohistochemical Protocol

Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP
Peroxidase/AP Blocker	5 min.	5 min.	5 min
Primary Antibody	30-60 min.	30-60 min.	30-60 min.
1st Step Detection	10 min.	30-45 min.	15 min.
2nd Step Detection	10 min.	Not Applicable	15 min.
Substrate-Chromogen	5-10 min.	5-10 min.	5-10 min.
Counterstain / Coverslip	Varies	Varies	Varies

Mounting Protocols

For detailed instructions using biodegradable permanent mounting media such as XyGreen PermaMounter (BSB 0169-0174) or organic solvent-based resin such as PermaMounter (BSB 0094-0097), refer to PI0174 or PI0097.

Product Limitations

Due to inherent variability present in immunohistochemical procedures (including fixation time of tissues, dilution factor of antibody, retrieval method utilized and incubation time), optimal performance should be established using positive and negative controls. Results should be interpreted by a qualified medical professional.

References

1. Huang EJ, Reichardt LF. Trk receptors: roles in neuronal signal transduction. *Annu. Rev. Biochem.* 2003; 72: 609-642.
2. Segal, RA. Selectivity in Neurotrophin Signaling: Theme and Variations. *Annual Review of Neuroscience.* 2003; 26: 299-330.
3. Barbacid M, et al. The trk family of tyrosine protein kinase receptors". *Biochim. Biophys. Acta.* 1991; 1072 (2-3): 115-127.
4. Vaishnavi A, Le AT, Doebele RC. TRKING down an old oncogene in a new era of targeted therapy. *Cancer Discov.* 2015; 5(1): 25-34.
5. Hechtman JF, et al. Pan-Trk Immunohistochemistry Is an Efficient and Reliable Screen for the Detection of NTRK Fusions. *Am J Surg Pathol.* 2017 Nov;41(11):1547-1551.
6. A. Amatu, et al. Tropomyosin receptor kinase (TRK) biology and the role of NTRK gene fusions in cancer. *Annals of Oncology* 2019; 30 (Supplement 8): 5-15.
7. Mauri G, Valtorta E, Cerea G et al. TRKA expression and NTRK1 gene copy number across solid tumours. *J Clin Pathol* 2018; 71(10): 926-931.
8. Yeh I, Tee MK, Botton T et al. NTRK3 kinase fusions in Spitz tumours. *J Pathol* 2016; 240(3): 282-290
9. U.S. Department of Health and Human Services: Centers for Disease Control and Prevention. Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories. Supplement / Vol. 61, January 6, 2012.

Symbol Key / Légende des symboles/Erläuterung der Symbole

	Storage Temperature Limites de température Zulässiger Temperaturbereich		Manufacturer Fabricant Hersteller	REF	Catalog Number Référence du catalogue Bestellnummer
	Read Instructions for Use Consulter les instructions d'utilisation Gebrauchsanweisung beachten		Expiration Date Utiliser jusque Verwendbar bis	LOT	Lot Number Code du lot Chargenbezeichnung

