


OX-40/CD134

Clone: BSB-90

Mouse Monoclonal

**ASR**
Bio SB
 BIOSCIENCE FOR THE WORLD

www.biosb.com

Inset: IHC of OX-40/CD134 on a FFPE Lymph Node Tissue

Intended Use

Analyte Specific Reagent.

Analytical and performance characteristics for OX-40 antibody, clone BSB-90 are not established.

Immunogen

Synthetic peptide corresponding to the N-terminus of the human OX40 protein.

Summary and Explanation

OX-40 also known as CD134 and Tumor Necrosis Factor Receptor Superfamily Member 4 (TNFRSF4), is a member of the TNFR-superfamily of receptors which is not constitutively expressed on activated CD4 and CD8 T cells as well as a number of other lymphoid and non-lymphoid cells, unlike CD28. OX40 and its binding partner, OX40L (CD252), are costimulatory signals from OX40 to a conventional T cell to promote division and survival, augmenting the clonal expansion of effector and memory populations as they are being generated to antigen. OX40 and OX40L also regulate cytokine production from T cells, antigen-presenting cells, NK cells, and NKT cells, and modulate cytokine receptor signaling. In line with these important modulatory functions, OX40/OX40L interactions have been found to play a central role in the development of multiple inflammatory and autoimmune diseases. Besides the essential role played by OX40 signaling in generating memory CD4 T cells, recent reports show that it also has a unique role in generating memory CD8 T cells. In addition, recent genome-wide association studies have identified single-nucleotide polymorphisms of the OX40L and OX40 genes that are related to cardiovascular diseases and SLE, providing direct evidence for the involvement of the OX40-OX40L interaction in human diseases.

OX40 is a potent costimulatory receptor that can potentiate T-cell receptor signaling on the surface of T lymphocytes, leading to their activation by a specifically recognized antigen. In particular, OX40 engagement by ligands present on dendritic cells dramatically increases the proliferation, effector function, and survival of T cells. Preclinical studies have shown that OX40 agonists increase antitumor immunity and improve tumor-free survival by increasing T and B cell responses to reporter antigen immunizations, led to preferential upregulation of OX40 on CD4(+) FoxP3(+) regulatory T cells in tumor-infiltrating lymphocytes, and increased the antitumor reactivity of T and B cells in patients with melanoma.

Antibody Type	Mouse Monoclonal	Clone	BSB-90
Isotype	IgG1	Reactivity	Paraffin, Frozen
Localization	Cytoplasmic, Membranous	Control	Tonsil, Lymph Node, Thymus
Species Reactivity		Human	

Presentation

OX-40/CD134 is a mouse monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH 7.5, containing BSA and sodium azide as a preservative.

Catalog No.	Antibody Type	Dilution	Volume/Qty
BSB 3120	Tinto Prediluted	Ready-to-Use	3.0 mL
BSB 3121	Tinto Prediluted	Ready-to-Use	7.0 mL
BSB 3122	Tinto Prediluted	Ready-to-Use	15.0 mL
BSB 3123	Concentrated	1:25 - 1:100	0.1 mL
BSB 3124	Concentrated	1:25 - 1:100	0.5 mL
BSB 3125	Concentrated	1:25 - 1:100	1.0 mL

Control Slides Available

Catalog No.	Quantity
BSB 3126	5 slides

Precautions

1. For professional users only. Results should be interpreted by a qualified medical professional.
2. This product contains <0.1% sodium azide (NaN₃) as a preservative. Ensure proper handling procedures are used with this reagent.
3. Always wear personal protective equipment such as laboratory coat, goggles and gloves when handling reagents.
4. Dispose of unused solution with copious amount of water.
5. Do not ingest reagent. If reagent is ingested, seek medical advice immediately.
6. Avoid contact with eyes. If contact occurs, flush with large quantities of water.
7. Follow safety precautions of the heating device used for epitope retrieval (TintoRetriever Pressure Cooker or similar).
8. For additional safety information refer to Safety Data Sheet for this product.
9. For complete recommendations for handling biological specimens, please refer to the CDC document, "Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories" (see References in this document).

Storage Store at 2-8°C

Stability

This product is stable up to the expiration date on the product label. Do not use after expiration date listed on package label. Temperature fluctuations should be avoided. Store appropriately when not in use, and avoid prolonged exposure to room temperature conditions.

Specimen Preparation

Paraffin sections: The antibody can be used on formalin-fixed paraffin-embedded (FFPE) tissue sections. Ensure tissue undergoes appropriate fixation for best results. Pre-treatment of tissues with heat-induced epitope retrieval (HIER) is recommended using Bio SB ImmunoDNA Retriever with Citrate (BSB 0020-BSB 0023), ImmunoDNA Retriever with EDTA (BSB 0030-BSB 0033) or ImmunoDNA Digestor (BSB 0108-0112). See reverse side for complete protocol. Tissue should remain hydrated via use of Bio SB Immuno/DNA Washer solutions (BSB 0029 & BSB 0042).

Frozen sections and cell preparations: The antibody can be used for labeling acetone-fixed frozen sections and acetone-fixed cell preparations.

This Antibody has been quality control tested by immunohistochemistry as follows

Quality Control Procedure

Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP
Peroxidase/AP Blocker	5 min.	5 min.	5 min
Primary Antibody	30-60 min.	30-60 min.	30-60 min.
1st Step Detection	10 min.	30-45 min.	15 min.
2nd Step Detection	10 min.	Not Applicable	15 min.
Substrate-Chromogen	5-10 min.	5-10 min.	5-10 min.
Counterstain / Coverslip	Varies	Varies	Varies

Mounting Protocols

For detailed instructions using biodegradable permanent mounting media such as XyGreen PermaMounter (BSB 0169-0174) or organic solvent based resin such as PermaMounter (BSB 0094-0097), refer to PI0174 or PI0097.

Product Limitations

Due to inherent variability present in immunohistochemical procedures (including fixation time of tissues, dilution factor of antibody, retrieval method utilized and incubation time), optimal performance should be established through the use of positive and negative controls. Results should be interpreted by a qualified medical professional.

References

- Kawamata S, Hori T, Imura A, Takaori-Kondo A, Uchiyama T. Activation of OX40 signal transduction pathways leads to tumor necrosis factor receptor-associated factor (TRAF) 2- and TRAF5-mediated NF-kappaB activation. *J. Biol. Chem.* 1998; 273 (10): 5808-14.
- Michael Croft, et al. The Significance of OX40 and OX40L to T cell Biology and Immune Disease. *Immunol Rev.* 2009; May; 229(1): 173-191.
- Ishii N, et al. OX40-OX40 ligand interaction in T-cell-mediated immunity and immunopathology. *Adv Immunol.* 2010; 105:63-98.
- Curti BD, et al. OX40 is a potent immune-stimulating target in late-stage cancer patients. *Cancer Res.* 2013 Dec 15;73(24):7189-98.
- U.S. Department of Health and Human Services: Centers for Disease Control and Prevention. Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories. Supplement / Vol. 61, January 6, 2012.

Symbol Key / Légende des symboles/Erläuterung der Symbole

	Storage Temperature Limites de température Zulässiger Temperaturbereich		Manufacturer Fabricant Hersteller	REF	Catalog Number Référence du catalogue Bestellnummer
	Read Instructions for Use Consulter les instructions d'utilisation Gebrauchsanweisung beachten		Expiration Date Utiliser jusque Verwendbar bis	LOT	Lot Number Code du lot Chargenbezeichnung



69 Santa Felicia Dr., Santa Barbara, CA 93117, USA
Tel. (805) 692-2768 | Tel. (800) 561-1145 | Fax. (805) 692-2769
E-mail: sales@biosb.com | Website: www.biosb.com